

<http://dx.doi.org/10.11646/zootaxa.3793.4.8><http://zoobank.org/urn:lsid:zoobank.org/pub:D85192FC-EB6A-4EBD-A17E-F78BB9F717BA>***Gobduntaulepis*, a new generic name for *Acantholepis* Pianovskij, 1989
(Conodonta) non Krøyer, 1846 (Actinopterygii)**

ALEXANDER B. DOWELD

National Institute of Carpology (Gaertnerian Institution), 21 Konenkova Street, RUS-127560, Moscow, Russian Federation.

E-mail: nicar-sekretariat@yandex.ru; sekretariat@doweld.pro

The fossil genus *Acantholepis* (type species *A. bispinatus* Pianovskij, 1989, by original designation) was established by Pianovskij (in Pianovskij, Pianovskaja & Aleksandrova, 1989: 69) for a presumable fossil conodont from the Ordovician of Southern Tien Shan (Gobduntau mountains, Nuratau ridge, Southern Tien Shan, Uzbekistan). However, several years earlier, the identical generic names *Acantholepis* Krøyer, 1846: 98 for a living fish in current use (Actinopterygii: Osmeriformes: Argentinidae), *Acantholepis* Mayr, 1861: 26 for a genus of living ants (Formicidae) and *Acantholepis* Newberry, 1875: 38 for a distinctive fossil fish genus, both latter replaced since then by new generic names, all preceded the recently published conodont generic name, which is therefore preoccupied, having three senior homonyms.

Since the fossil conodont genus is recognized taxonomically as a valid genus, not being merged into another one, and its name does not have any senior or junior synonym, that could be used as a substitute name, a *nomen novum*, *Gobduntaulepis* nom. nov. (type species *Acantholepis bispinatus* Pianovskij, 1989), is here proposed in accordance with the International Code of Zoological Nomenclature, as a new replacement name for *Acantholepis* Pianovskij, 1989.

Systematics**Phylum Chordata Hackel, 1874****Class Conodonta Eichenberg, 1930**

Genus ***Gobduntaulepis***, *nomen novum pro Acantholepis* Pianovskij in Pianovskij, Pianovskaja & Aleksandrova, 1989: 69.

Etymology. The name Gobduntau (G‘ubdintog‘ in Uzbek) is that of the mountains (39°55’00” N, 67°15’00” E) which are the type locality of the type species of this nominal genus.

Grammatical gender: masculine.

Type species. *Gobduntaulepis bispinatus* (Pianovskij in Pianovskij, Pianovskaja & Aleksandrova, 1989: 69) through original designation under *Acantholepis* Pianovskij in Pianovskij, Pianovskaja & Aleksandrova, 1989: 69.

Other species. *Gobduntaulepis dubius* (Pianovskij in Pianovskij, Pianovskaja & Aleksandrova, 1989: 69) comb. nov.; *Gobduntaulepis exolimbus* (Pianovskij in Pianovskij, Pianovskaja & Aleksandrova, 1989: 70) comb. nov. *Stratum typicum* of the type species: Lower Ordovician (Dzhalatarian suite [lower part]) (i.e., Tremadoc-Arenig).

Geography. Kattasaj, Gobduntau mountains & Narvan saj basin, Northern Nuratau ridge, Southern Tien Shan, Uzbekistan.

References

Eichenberg, W. (1930) Conodonten aus dem Culm des Harzes. *Palontologische Zeitschrift*, 12, 177–182.

<http://dx.doi.org/10.1007/bf03044446>

Hackel, E. (1874) *Anthropogenie oder Entwicklungsgeschichte des Menschen*. W. Engelmann, Leipzig, Tafeln, 210 Textabb. und 36 Tab. 1 Bl., XVI S., 1 Bl., 732 S.

- Krøyer, H.N. (1846) *Danmarks Fiske. Vol. 3*. S. Triers, Copenhagen, 1279 pp.
- Mayr, G.L. (1861) *Die Europäischen Formiciden: nach der analytischen Methode bearbeitet*. C. Gerold's Sohn, Wien, 44 pp.
- Newberry, J.S. (1875) Description of fossil fishes. *Report of the Geological Survey of Ohio*, 2 (2), 1–64.
- Pianovskij, G.V., Pianovskaja, I.A. & Aleksandrova, N.V. (1989) Probable conodont animals from Predevonian deposits of Southern Tien Shan (western part) and their associated fauna]. *In: Pianovskaja, I.A. (Ed.), Biostratigraphic Researches in Uzbekistan*. SAIGIMS, Tashkent, pp. 54–72. [Пяновский, Г.В., Пяновская, И.А., Александрова, Н.В. (1989) Вероятные конодонтовые животные из додевонских отложений южного Тянь-Шаня (западная часть) и ассоциирующаяся с ними фауна. *In: Пяновская, И.А. (Ed.), Биостратиграфические исследования в Узбекистане*] [in Russian]